

A Sea Anemone Outbreak Eliminates Damselfish Territories from Fringing Reefs in Southern Taiwan

Tsen-Chien Chen¹, Cheng-Tze Ho², and Rong-Quen Jan^{2*}

¹ *Department of Leisure Management, Minghsin University of Science and Technology, Hsinchu 30401, Taiwan*

² *Research Center for Biodiversity, Academia Sinica, Taipei 11529, Taiwan*

(Accepted February 26, 2008)

In 2000, when the sea anemone, *Mesactinia ganesis*, was rare on fringing reefs in an embayment in southern Taiwan (21°57'N; 120°45'E) (Jan et al. 2007), bluntnout gregory *Stegastes lividus* (Bloch and Schneider) (Fig. 1a) and dusky gregory, *S. nigricans* (Lacepède) (Jan et al. 2003) had long-established territories on coral branches on the reef flat. An outbreak of the sea anemone has occurred in recent years in this embayment. Colonies of branching corals on the slanting reef surface were eliminated by the spreading *M. ganesis*. Strong territoriality of the damselfishes has slowed down the elimination process, as remaining coral colonies can still be found within their territories (Fig. 1b).




Fig. 1. Transitions of territorial changes of *Stegastes lividus*. (a) An adult and a patch of an algal mat growing on dead branches of the coral *Acopora muricata* as the focal point of its territory; (b) an adult and its territory surrounded by sea anemone colonies; (c) coral colonies shared by a group of *S. lividus* and *S. nigricans*. Note that the reef substratum is mostly covered by sea anemone colonies. The arrow indicates the territory holder.

Nevertheless, currently more than 80% of the former damselfish territories have already disappeared from this part of the reef (Fig. 2). Some holders emigrated to the lower part of a near-by reef, competing for new territories on the few subsisting

coral colonies (Fig.1c). In parallel, relic coral skeletons from abandoned damselfish territories have been taken over by Dick's damsel, *Plectroglyphidodon dickii* (Liénard), as nesting substrate. This observation indicates that the sea anemone outbreak has resulted in drastic changes in the demography of habitat specialist in fishes. While the outbreak has shown no signs of recession, the fish community, along with other fouling organisms, will likely remain unstable in this embayment.

<http://zoolstud.sinica.edu.tw/Journals/47.3/xxx.pdf>




Fig. 2. Yearly changes in the number of *Stegastes* territories (mean for 4 seasonal counts, with the standard deviation; filled columns: *S. lividus*; empty columns: *S. nigricans*) built on coral colonies in a fixed quadrat of 45 m².

Acknowledgments: This work was supported by Long-Term Ecological Research (NSC92-2621-B-001-003 and NSC93-2621-B-001-005) and NSC96-2621-B-001-009-MY3 from the National Science Council of Taiwan and a competitive grant to RQJ from the Research Center for Biodiversity, Academia Sinica.

REFERENCES

Jan RQ, CT Ho, FK Shiah. 2003. Determinants of territory size of the dusky gregory. *J. Fish. Biol.* **63**: 1589-1597.

Jan RQ, YT Shao, FP Lin, TY Fan, YY Tu, HS Tsai, KT Shao. 2007. An underwater camera system for real-time coral reef fish monitoring. *Raffles Bull. Zool.* **14(Supplement)**: 273-279.

*To whom correspondence and reprint requests should be addressed. °Tel: 886-2-27899519. E-mail:rqjan@gate.sinica.edu.tw